

Durham Research Online

Deposited in DRO:

05 April 2017

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Paschalidou, A.K. and Kassomenos, P.A. and McGregor, G.R. (2017) 'Analysis of the synoptic winter mortality climatology in five regions of England : searching for evidence of weather signals.', *Science of the total environment.*, 598 . pp. 432-444.

Further information on publisher's website:

<https://doi.org/10.1016/j.scitotenv.2017.03.276>

Publisher's copyright statement:

© 2017 This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Analysis of the synoptic winter mortality climatology in five regions of England: Searching for evidence of weather signals

A.K. Paschalidou^{1*}, P.A. Kassomenos² and G.R. McGregor³

¹Department of Forestry and Management of the Environment and Natural Resources,
Democritus University of Thrace, GR-68200, Greece

²Laboratory of Meteorology, Department of Physics, University of Ioannina, GR-45110,
Greece

³Department of Geography, Durham University, Durham, United Kingdom

* E-mail address: apascha@fmenr.duth.gr

Abstract: Although heat-related mortality has received considerable research attention, the impact of cold weather on public health is less well-developed, probably due to the fact that physiological responses to cold weather can vary substantially among individuals, age groups, diseases etc, depending on a number of behavioral and physiological factors. In the current work we use the classification techniques provided by the COST-733 software to link synoptic circulation patterns with excess cold-related mortality in 5 regions of England. We conclude that, regardless of the classification scheme used, the most oppressive conditions for public health in England are associated with the prevalence of the Easterly type of weather, favoring advection of cold air from continental Europe. It is noteworthy that there has been observed little-to-no regional variation with regards to the classification results among the 5 regions, suggestive of a spatially homogenous response of mortality to the atmospheric patterns identified. In general, the 10 different groupings of days used reveal that excess winter mortality is linked with the lowest daily minimum/maximum temperatures in the area. However it is not uncommon to observe high mortality rates during days with higher, in relative terms, temperatures, when rapidly changing weather results in an increase of mortality. Such a finding confirms the complexity of cold-related mortality and highlights the importance of synoptic climatology in understanding of the phenomenon.

Keywords: Cold-related mortality, classification methods, atmospheric circulation, Easterly weather

1. Introduction

Extreme weather, in the form of heat waves or cold spells, is associated with adverse health effects in many regions of the world, where strong links between ambient temperature and increased mortality have been reported (Ferreira Braga et al., 2001; Basu and Samet, 2002; Donaldson et al., 2003; Basu et al., 2008; Basu, 2009; Guo et al., 2013; Urban et al., 2014;

Wang et al., 2014; Tsangari et al., 2016). However, the exact shape of the exposure-response curve has been found to vary with location and latitude, depending on a number of physiological and behavioral factors (Guo et al., 2014; Keatinge et al., 2000). Physiological factors include acclimatization to extreme ambient temperatures, as well as the ability of thermoregulation, while behavioral factors include the habits and lifestyle of population, the use of air-conditioning, the quality of housing, the time spent outdoors etc (Donaldson and Keatinge, 2003; Kovats and Kosatsky, 2009; Yu et al., 2012).

According to Carson et al. (2006) and Astrom et al. (2013) a progressive reduction in temperature-related mortality has been reported in many regions of the world since the beginning of the twentieth century. Nevertheless, extreme ambient temperatures still pose a severe threat to public health (Morabito et al., 2012; Scarborough et al., 2012; Astrom et al., 2013), especially for the most vulnerable groups of population, such as the children, the pregnant women and the elderly (Hajat et al., 2007; Xu et al., 2013). Under the changing climate, the frequency, intensity and duration of heat waves are expected to increase, resulting in an increase of heat-related mortality (Huang et al., 2011; Hajat et al., 2014; Heaviside et al., 2016), whereas winter mortality is expected to decrease. However, the future of winter mortality is not completely understood (Wang et al., 2016).

Cold spells have been long-known for their adverse health effects and their impact on mortality (Keatinge 2002). For instance, Curriero et al. (2002) have studied the sensitivity of population to cold weather in 11 cities in the United States, whereas Analitis et al. (2008) have studied the impact of cold spells on public health (in the form of cardiovascular, respiratory and cerebrovascular diseases) in 15 cities in Europe. In China, a severe cold spell in 2008 resulted in 148279 excess deaths, with the highest impact observed in southern and central China, where mortality increased by 44% (Ma et al., 2013; Xie et al. 2013; Zhou et al., 2014). Similarly, the adverse health effects of cold weather have been studied in many areas of the Iberian Peninsula (Gomez-Acebo et al., 2010, Montero et al., 2010; Vasconcelos et al., 2013) and Italy, where an unusual cold spell during 2012 resulted in a 25% increase in mortality among the elderly (75+ years of age) across 14 cities (de' Donato et al., 2013). In England, winter temperature seasonality and cold spells linked to a severe negative phase of the North Atlantic Oscillation have been associated with increased risk of ischemic heart disease and myocardial infarction (McGregor 2005; Bhaskaran et al., 2010).

The majority of studies described above use time series-based Poisson Regression and Generalized Additive Models (GAMs) and/or case-only or crossover studies to analyze the cold weather-related mortality. However, the climatology of winter mortality is less well-developed, mainly due to the fact that winter mortality is very often confounded by social, economic, behavioral and physiological factors (Anderson and Bell 2009; Allen and Lee, 2014). The objective of this paper is to shed light on the climatological associations between winter mortality and cold weather in the United Kingdom, by using synoptic classifications, and to explore the possible link between certain atmospheric patterns and winter mortality, so that the most oppressive conditions for public health can be recognized.

2. Data and Methods

2.1 Area Description and Data Sources

In the present study we focus on the United Kingdom, where cold spells are well-known to play an important role in the yearly variability of excess cold-related mortality (McGregor 2005; Hajat and Kovats 2014). According to Healy (2003), the country presents the highest levels of excess cold-related mortality across Europe, with the cold-related mortality burden accounting for more than one order of magnitude more deaths than heat-related mortality (approximately 61 and 3 deaths per 100,000 population per year, respectively) (Vardoulakis et al., 2014).

For the needs of the study, November to February daily temperature (surface mean, minimum and maximum, in °C) and mortality (all-cause deaths per day) data for the 26-yr period 1974-1999 were used. The data covered the following 5 official Office of National Statistics (ONS) regions in England: (a) Yorkshire and the Humber, (b) the West Midlands, (c) Northeast, (d) Northwest and (e) Southeast regions (Figure 1), as these were the only regions for which mortality data could be obtained. The above regions can be considered to represent the whole of England, as they capture the range of winter temperatures across England, and could therefore provide insights into any geographical variation of cold weather and adverse health effects.

It is noted that the temperature data were obtained from one county-level meteorological station, representative of each region (Figure 1), namely, West Yorkshire (Yorkshire and the Humber), West Midlands (West Midlands), Tyne and Wear (Northeast), Greater Manchester (Northwest), and Hampshire (Southeast). Inevitably small temperature differences probably appeared among the different areas of each region. Nevertheless, the use of the county-level meteorological stations for representing each region is probably the best option, under the circumstances, as no additional information is available on a number of important issues, such as whether the people travelled within the region (or even throughout the country) before dying, in which part of the region they resided etc.

All data were obtained from the ONS and the mortality data were de-trended prior to the analysis.

2.2 Methodology

In order to link winter mortality and prevailing weather systems, the COST-733 Action (<http://www.cost733.eu>) classification tool (v. 2.0) was used. This software uses a number of well-established classification methods to provide synoptic classification schemes for Europe. For the purposes of the present study, Domain 4 of this software, including 432 cells defined by the 47-62°N and 18-8°E coordinates, and covering the British Isles, Benelux and N. France area was used. For the needs of the study the following 10 available grouping techniques were used: (a) 2 schemes using the Leader Algorithm, namely the ERP (based on the pressure gradient metric; Ericum et al., 2008) and the LND (based on the correlation coefficient metric; Lund, 1963), (b) 4 Principal Component Analysis (PCA) schemes, namely the KRZ (S-mode without rotation), the PCT (T-mode with oblique rotation), the PTT (T-mode with VARIMAX rotation) and the PXE (S-mode with VARIMAX rotation) and (c) 4 Optimization

Algorithms, namely the CAP (principle components cluster analysis), the CKM which is a k-means by dissimilar seeds, the PXX (K-means reassigned extreme scores) and the SAN (simulated annealing and diversified randomization clustering).

The Leader Algorithm counts the number of elements with similarity to the key-pattern exceeding a certain threshold, in order to find the so-called “leader”, i.e. representative key-patterns for each group. A T-mode PCA recognizes typical patterns and uses loadings to describe the degree of realization, while an S-mode PCA locates temporal variability through typical modes. Finally, the Optimization Algorithms are non-hierarchical methods, in the sense that the various groups can be split up to minimize the within-group variance. A complete description of the aforementioned classification schemes, together with their advantages and disadvantages can be found in Philipp et al. (2010, 2014).



Figure 1. Office of National Statistics study regions.

The grouping techniques described above were used to associate synoptic circulations and winter mortality and to investigate any evidence of weather signals. The groupings were carried out with a 4 days sequence length (i.e. a 4-days averaging), using mean sea-level pressure as the classification variable. The 4 days sequence length was used, as winter mortality can manifest itself in a time distributed way, often referred to as the “lag effect” (Gasparini et al., 2010; Allen and Sheridan, 2014; Zeka et al., 2014), as opposed to heat-related mortality which has an acute response (Gosling et al., 2009). The various classification schemes used resulted in a number of classes (9 or 10 depending on the specific configuration used) with distinct synoptic circulation patterns. Then, mortality data were summarized across the different synoptic circulations.

Next, we used the PI sign-test (Duckstein et al., 1993; Paschalidou and Kassomenos, 2016) to check the frequency of excess mortality in the various classes:

$$PI_i = 100 \times \left(\frac{\text{number of deaths in } C_i / \text{total number of deaths}}{\text{number of days in } C_i / \text{total number of days}} - 1 \right) \quad (1)$$

where C_i stands for the different classes. In Eq. 1, values of PI equal to 0 denote mortality is equally spread among classes, whereas positive/negative PI values show that the occurrence of deaths is more/less frequent under the conditions of the specific classes. PI values equal to -100 indicate “mortality-free” classes.

Finally, centroid maps produced by the COST-733 software were discussed, in order to study the synoptic winter mortality climatology.

3. Results and Discussion

3.1 Presentation of the classes

During the 26-y period 1974-1999 a total of 2916300 all-cause casualties were recorded in the 5 regions studied. The number of deaths per year for each of the five regions is presented in Figure 2. Not surprisingly the biggest number of fatalities for most of the years appears in the Northwest region (194.6 deaths per year), where, according to the Office of National Statistics, this is the most populated region (approximately 498 inhabitants per Km^2). A closer look reveals a progressive reduction in winter mortality for all five regions of England. Although not all of the incidents can be attributed/related to cold weather, the general decreasing trend is in agreement with Carson et al. (2006) and Astrom et al. (2013) who reported descending trends in cold-related mortality for London and Stockholm, respectively. Such a decreasing trend probably reflects the changing vulnerability of the population due to improvements in infrastructure, lifestyle, technology, and general health. According to Vardoulakis et al. (2014), the decreasing trend is projected to continue due to climate change and to reach approximately 42 deaths per 100,000 population per year in the

UK, whereas the heat-related mortality burden is expected to increase to approximately 9 deaths per 100,000 population per year by the 2080s.




Figure 2. The number of deaths per year for each of the five regions

Tables 1-5 present the results of all 10 classification techniques performed for each one of the 5 regions studied, namely the number of days per class, the maximum/minimum/mean surface temperature (in °C) and the total number of deaths. It appears that for each classification scheme there are a number of classes that reflect the most oppressive conditions, in terms of excess winter mortality. This finding is highlighted in Figure 3, where the PI index for each classification scheme is shown for the Northwest region (figures for the rest of the regions are omitted).

Specifically for the Northwest region ERP classes C6 and C8 reflect the most oppressive conditions for public health, as they present the biggest PI values (Figure 3). It is noteworthy that these classes feature the lowest maximum (4.7 °C and 4.5 °C, respectively), as well as the lowest minimum temperatures (0.3 °C and -2.3 °C, respectively), confirming that winter mortality is associated with low temperatures (Keatinge 2002). Similar is the pattern for the Southeast (Table 2), the West Midlands (Table 3), the Yorkshire and Humber (Table 4) and the Northeast region (Table 5), where classes C6 and C8 presenting the lowest maximum/minimum temperatures are linked with the highest PI values (figures not shown). This finding is in agreement with results provided by Dimitriou et al. (2016) who defined atmospheric pathways linked with winter low temperature episodes (LTE) in the same 5 regions of England for the same time period and revealed associations with excess mortality rates. According to them, a statistically significant increase in mortality was calculated for LTE days across all 5 regions studied. LND class C3, with PI value equal to 9.04, appears to be the most dangerous for public health in the Northwest region. This class comprises 10% of the total winter days (November – February) for the period studied and is linked with the


lowest maximum and minimum temperatures in the region (4.6 °C and 0.4 °C, respectively). The pattern of the highest mortality being linked to the lowest temperatures is repeated in the Southeast (Table 2), the West Midlands (Table 3), the Yorkshire and Humber (Table 4) and the Northeast region (Table 5).

Among the 4 PCA classifications schemes in the Northwest region, namely KRZ, PCT, PTT and PXE, KRZ class C7, PCT class C8, PTT class C3 and PXE class C9 (Figure 3) appear to be linked with the most adverse effects on public health. These classes comprise 10%, 9%, 20% and 8%, respectively, of the total winter days studied. Although KRZ class C7 presents some of the lowest maximum and minimum temperatures (5.5 °C and 1.0 °C, respectively), classes PCT C8, PTT C3, PXE C9 are associated with slightly higher maximum (6.7 °C, 6.9 °C and 7.8 °C, respectively) and minimum temperatures (2.4 °C, 2.1 °C and 3.3 °C, respectively). Similar is the situation in the Southeast (Table 2), the West Midlands (Table 3), the Yorkshire and Humber (Table 4) and the Northeast region (Table 5), where the highest levels of winter mortality are not always linked to the lowest temperatures. This finding does not come as a surprise, as Hajat and Kovats (2014) state that a significant number of cold-related deaths are linked to moderate temperatures. Additionally, Gasparrini and Leone (2014) conclude that 70% of cold-related mortality in London is observed during days with temperature higher than 5 °C and not necessarily during days with the absolute minimum temperatures. Such a pattern could also mean there is a temperature range (zone), centered around the lowest temperatures, that could cause the excess mortality. In the presence of various confounding factors, it can be the slightly higher temperatures that are linked to excess mortality. Furthermore, Dimitriou et al. (2016) concluded that in some cases mortality in the 5 regions studied is linked to rather increased winter temperatures associated with long-distance west-to-east flows, resulting in rapidly changing weather, as opposed to stable conditions linked with blocking to the east. Rapidly changing atmospheric pressure and/or temperature have been longed blamed for adverse health effects. For instance Dawson et al. (2008) showed that rapid falls in atmospheric pressure are associated with increased risk of morbidity due to hemorrhagic stroke in Glasgow. Finally, the excess mortality during days with relatively higher temperatures could be explained as lagged mortality after a period of cumulative exposure to oppressive cold weather.

With regards to the 4 optimization algorithm classification schemes, CAP class C2, CKM class C2, PKX class C10 and SAN class C4 are associated with the most dangerous conditions for public health, in terms of winter mortality in the North west region. These classes comprise 8%, 10%, 6% and 9%, respectively, of the total winter days studied and present some of the lowest maximum (6.1 °C, 5.1°C, 4.2°C and 5.9°C, respectively) and minimum temperatures (1.3 °C, 0.3°C, 0.0°C and 1.2 °C, respectively) in the region (Table 1). Almost identical is the pattern observed for the Southeast (Table 2), the West Midlands (Table 3), the Yorkshire and Humber (Table 4) and the Northeast region (Table 5), where CKM C2 and PXE C10 are associated with the lowest temperatures and the highest levels of excess mortality, while classes CAP C2 and SAN C4 also present some of the lowest maximum/minimum temperatures.


On the whole, the pattern of the “most dangerous” classes described above is repeated almost identically (Tables 2-5), confirming the adverse impact of the specific classes’

climatology on winter mortality. Specifically, there appears to be little-to-no regional variation with regards to the most oppressive classes among the 5 regions, suggesting a spatially homogeneous response of mortality in the 5 regions to atmospheric patterns.



(a)

(b)



(c)

(d)




Figure 3. The PI index together with the minimum and maximum temperature for the 10 classification schemes, in the Northwest region

Table 1. The number of days for each class, together with the maximum/minimum/mean temperature (in °C) and the total mortality for each classification scheme in the Northwest region. The classes showing the most oppressive (in terms of excess mortality) conditions are given in bold

		CLASSES									
		1	2	3	4	5	6	7	8	9	10
ERP	Total days	168	628	595	499	765	223	219	21	8	-
	Max T	6.5	6.4	9.4	6.7	9.1	4.7	8.1	4.5	9.6	-
	Min T	1.7	1.4	4.2	1.1	3.6	0.3	2.3	-2.3	3.4	-
	Mean T	3.9	3.9	6.7	3.8	6.3	2.5	5.2	1.1	6.5	-
	Tot Mort	31947	124778	111856	93629	149910	47095	43041	4363	1609	-
LND	Total days	680	472	304	439	262	210	330	209	220	-
	Max T	9.6	6.8	4.6	9.0	7.3	6.1	8.1	8.2	6.4	-
	Min T	4.2	1.1	0.4	3.4	2.1	0.7	3.0	2.9	1.6	-
	Mean T	6.8	3.8	2.5	6.1	4.7	3.3	5.4	5.4	3.9	-
	Tot Mort	129048	90018	64493	85280	50174	38938	65891	40257	44129	-
KRZ	Total days	617	427	564	291	180	275	324	222	226	-
	Max T	9.5	9.6	8.9	7.0	6.9	6.3	5.5	5.3	4.9	-
	Min T	4.0	3.9	3.5	2.1	1.8	0.7	1.0	0.6	-0.5	-
	Mean T	6.6	6.7	6.1	4.5	4.3	3.5	3.2	2.9	2.1	-
	Tot Mort	119727	80782	106438	58426	34948	52308	67312	44105	44183	-
PCT	Total days	503	554	237	433	343	298	207	294	257	-
	Max T	8.2	9.9	5.2	8.7	5.9	7.4	7.4	6.7	6.6	-
	Min T	2.5	4.5	-0.0	3.0	1.3	2.3	2.1	2.4	1.0	-
	Mean T	5.3	7.1	2.5	5.8	3.5	4.7	4.7	4.5	3.8	-
	Tot Mort	95618	105781	45797	84069	69608	56364	40329	60680	49983	-
PTT	Total days	1129	916	612	195	87	65	59	19	44	-
	Max T	8.9	7.9	6.9	5.9	5.7	4.4	5.5	5.7	6.2	-
	Min T	3.5	2.3	2.1	0.1	1.5	-0.7	0.2	1.9	1.7	-
	Mean T	6.1	5.0	4.4	3.0	3.5	1.8	2.9	3.7	3.9	-
	Tot Mort	216301	173800	125246	38556	17281	13224	11746	3667	8406	-
PXE	Total days	551	525	346	238	301	329	217	185	235	199
	Max T	9.1	9.0	8.2	7.5	8.9	6.4	5.1	4.7	7.8	5.6
	Min T	3.6	3.6	2.7	2.2	3.2	1.5	0.6	-0.9	3.3	0.2
	Mean T	6.2	6.2	5.4	4.8	6.0	3.9	2.8	1.9	5.5	2.8
	Tot Mort	107381	99379	65435	44911	57300	66602	44163	36497	49106	37453
CAP	Total days	304	236	297	388	504	391	306	245	455	-
	Max T	4.9	6.1	7.4	8.9	9.9	6.6	7.4	7.4	8.4	-
	Min T	-0.2	1.3	1.7	3.7	4.3	1.7	1.8	2.8	2.9	-
	Mean T	2.3	3.6	4.4	6.2	7.0	4.1	4.5	5.0	5.6	-
	Tot Mort	60566	49194	56365	74291	94865	74240	57921	49670	91117	-
CKM	Total days	480	303	392	437	228	244	338	398	306	-
	Max T	9.8	5.1	8.6	5.9	6.2	6.8	7.8	9.4	7.8	-
	Min T	4.2	0.3	3.5	0.9	0.4	2.3	2.1	3.8	2.7	-
	Mean T	6.9	2.6	5.9	3.4	3.3	4.5	4.9	6.5	5.2	-
	Tot Mort	91837	62454	74400	83732	43103	50026	63817	75999	62861	-
PXK	Total days	524	576	361	315	294	240	279	216	134	187

	Max T	9.2	9.6	6.9	5.9	7.8	5.9	9.1	7.3	5.5	4.2
	Min T	3.9	3.8	1.6	1.3	2.5	-0.1	4.1	2.2	0.4	0.0
	Mean T	6.5	6.6	4.2	3.5	5.1	2.9	6.5	4.6	2.9	2.1
	Tot Mort	100649	110011	67851	64430	57054	46569	56638	40252	25910	38866
SAN	Total days	267	278	435	269	481	397	241	369	389	-
	Max T	5.0	6.8	9.1	5.9	9.9	6.5	6.8	8.3	8.3	-
	Min T	-0.1	0.9	3.8	1.2	4.3	1.5	2.3	2.5	3.1	-
	Mean T	2.4	3.9	6.3	3.5	7.0	4.0	4.5	5.3	5.6	-
	Tot Mort	52056	52509	82886	56561	90857	75553	48906	69807	79093	-

269

270

271 Table 2. The number of days for each class, together with the maximum/minimum/mean
272 temperature (in °C) and the total mortality for each classification scheme in the Southeast
273 region. The classes showing the most oppressive (in terms of excess mortality) conditions
274 are given in bold

		CLASSES									
		1	2	3	4	5	6	7	8	9	10
ERP	Total days	168	628	595	499	765	223	219	21	8	-
	Max T	6.7	7.3	9.7	7.8	9.9	5.8	9.3	5.2	9.9	-
	Min T	1.3	1.6	4.1	1.2	4.4	0.6	3.2	-1.7	4.8	-
	Mean T	4.0	4.5	6.9	4.5	7.1	3.2	6.2	1.8	7.4	-
	Tot Mort	29083	113117	101300	85760	135453	42677	39016	4041	1351	-
LND	Total days	680	472	304	439	262	210	330	209	220	-
	Max T	10.5	7.7	5.1	9.6	8.2	7.5	8.7	8.4	6.9	-
	Min T	4.9	1.1	0.1	3.8	2.3	1.2	3.5	2.3	1.7	-
	Mean T	7.7	4.4	2.6	6.7	5.2	4.4	6.1	5.4	4.4	-
	Tot Mort	116564	81544	59361	76712	45858	35848	59039	37060	39810	-
KRZ	Total days	617	427	564	291	180	275	324	222	226	-
	Max T	10.1	10.3	9.9	7.9	7.7	7.3	6.2	5.7	5.7	-
	Min T	4.7	4.6	3.7	2.4	1.7	0.7	1.1	0.2	-0.2	-
	Mean T	7.4	7.4	6.8	5.2	4.7	4.0	3.6	3.0	2.7	-
	Tot Mort	108337	73092	96225	52793	31596	47971	61142	40668	39975	-
PCT	Total days	503	554	237	433	343	298	207	294	257	-
	Max T	9.2	10.5	5.8	9.1	6.5	9.4	7.6	7.7	7.1	-
	Min T	2.7	5.0	0.0	3.6	1.3	3.4	1.5	2.8	0.6	-
	Mean T	5.9	7.8	2.9	6.3	3.9	6.4	4.6	5.3	3.8	-
	Tot Mort	85756	95384	41881	76030	63905	51719	36807	54511	45804	-
PTT	Total days	1129	916	612	195	87	65	59	19	44	-
	Max T	9.7	8.9	7.5	6.3	5.3	4.2	5.7	7.2	7.8	-
	Min T	4.1	2.6	2.5	-0.5	0.4	-1.2	-0.7	2.2	2.4	-
	Mean T	6.9	5.8	5.0	2.9	2.9	1.5	2.5	4.7	5.1	-
	Tot Mort	195843	157534	113137	35195	15898	12060	10967	3408	7756	-
PXE	Total days	551	525	346	238	301	329	217	185	235	199
	Max T	9.2	10.3	8.8	8.6	10.0	6.9	5.3	5.0	8.9	7.1
	Min T	3.6	4.5	2.8	2.5	3.8	1.4	0.2	-1.2	4.2	0.9
	Mean T	6.4	7.4	5.8	5.6	6.9	4.1	2.7	1.9	6.5	4.0
	Tot Mort	97191	90147	59714	41312	51378	59669	41000	33167	43995	34224

CAP	Total days	304	236	297	388	504	391	306	245	455	-
	Max T	5.1	6.6	8.1	9.3	10.7	6.6	8.8	8.7	9.9	-
	Min T	-0.7	1.5	1.6	3.6	5.1	0.9	2.4	3.7	4.2	-
	Mean T	2.2	4.1	4.9	6.5	7.9	3.7	5.6	6.2	7.1	-
	Tot Mort	55807	44633	50976	67652	86231	68440	52366	44297	81397	-
CKM	Total days	480	303	392	437	228	244	338	398	306	-
	Max T	10.2	5.6	8.9	5.9	7.1	8.1	9.1	10.5	9.5	-
	Min T	4.3	0.2	2.8	0.2	0.6	2.9	2.7	5.0	3.9	-
	Mean T	7.5	2.9	5.9	3.0	3.8	5.5	5.9	7.8	6.7	-
	Tot Mort	83487	57170	67766	77098	39309	44952	57822	68459	55735	-
PXK	Total days	524	576	361	315	294	240	279	216	134	187
	Max T	10.1	10.1	8.9	6.6	7.8	6.6	10.1	7.9	5.7	4.7
	Min T	4.6	4.4	2.5	1.5	1.6	-0.2	4.9	2.1	-0.1	-0.2
	Mean T	7.3	7.2	5.7	4.1	4.7	3.2	7.6	5.0	2.8	2.2
	Tot Mort	91464	98332	61997	58484	52393	42294	50316	36931	23794	35792
SAN	Total days	267	278	435	269	481	397	241	369	389	-
	Max T	5.2	7.9	9.4	6.4	10.6	6.4	8.2	9.6	9.9	-
	Min T	-0.8	1.4	3.5	1.3	5.1	0.8	3.0	3.2	4.5	-
	Mean T	2.2	4.6	6.5	3.8	7.8	3.6	5.6	6.4	7.2	-
	Tot Mort	47851	47860	75383	51313	82461	69667	43936	62842	70485	-

275

276

277 Table 3. The number of days for each class, together with the maximum/minimum/mean
278 temperature (in °C) and the total mortality for each classification scheme in the West
279 Midlands region. The classes showing the most oppressive (in terms of excess mortality)
280 conditions are given in bold

		CLASSES									
		1	2	3	4	5	6	7	8	9	10
ERP	Total days	168	628	595	499	765	223	219	21	8	-
	Max T	6.0	5.9	9.2	6.7	9.0	4.1	8.4	3.7	9.5	-
	Min T	1.3	0.7	3.7	0.7	3.3	-0.7	2.4	-3.2	3.9	-
	Mean T	3.7	3.3	6.4	3.7	6.1	1.7	5.4	0.3	6.7	-
	Tot Mort	29845	116735	104550	88387	139961	44267	39737	4064	1445	-
LND	Total days	680	472	304	439	262	210	330	209	220	-
	Max T	9.8	6.7	3.7	8.8	7.3	6.2	7.5	7.6	5.6	-
	Min T	4.1	0.5	-0.8	2.9	1.8	0.6	2.3	1.8	0.8	-
	Mean T	7.0	3.6	1.4	5.9	4.6	3.4	4.9	4.7	3.2	-
	Tot Mort	119301	84370	61017	80150	47426	36725	60577	38265	41161	-
KRZ	Total days	617	427	564	291	180	275	324	222	226	-
	Max T	9.4	9.7	9.1	6.5	6.6	6.2	4.7	4.5	4.6	-
	Min T	3.7	3.7	3.2	1.4	1.0	0.1	0.1	-0.4	-1.1	-
	Mean T	6.6	6.7	6.1	4.0	3.8	3.2	2.4	2.1	1.7	-
	Tot Mort	111239	75406	99424	54524	32449	49479	63008	41766	41697	-
PCT	Total days	503	554	237	433	343	298	207	294	257	-
	Max T	8.2	9.9	4.8	8.5	5.0	8.1	7.0	6.1	6.2	-
	Min T	2.1	4.3	-0.5	2.5	0.3	2.3	1.3	1.7	0.0	-
	Mean T	5.2	7.1	2.1	5.5	2.7	5.2	4.2	3.9	3.1	-

	Tot Mort	88913	97377	43577	78502	65671	53097	37997	56526	47331	-
	Total days	1129	916	612	195	87	65	59	19	44	-
	Max T	8.9	7.9	6.1	5.4	4.8	3.5	4.9	5.6	6.4	-
PTT	Min T	3.3	1.9	1.3	-1.2	0.6	-1.5	-0.7	1.2	1.3	-
	Mean T	6.1	4.9	3.7	2.1	2.7	1.0	2.1	3.4	3.8	-
	Tot Mort	201008	163368	116871	36531	15871	12596	11175	3625	7946	-
	Total days	551	525	346	238	301	329	217	185	235	199
	Max T	8.5	9.4	8.2	7.7	9.1	5.7	4.0	4.3	7.4	5.6
PXE	Min T	2.8	3.5	2.3	1.9	2.9	0.5	-0.3	-1.6	2.8	0.1
	Mean T	5.7	6.4	5.2	4.9	6.0	3.1	1.8	1.3	5.1	2.8
	Tot Mort	100058	92345	60778	42617	53952	62290	41721	34375	45394	35461
	Total days	304	236	297	388	504	391	306	245	455	-
	Max T	4.1	5.3	7.3	8.8	10.1	5.9	7.6	6.9	8.6	-
CAP	Min T	-1.3	0.4	1.1	3.1	4.2	0.8	1.6	2.2	2.8	-
	Mean T	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	-
	Tot Mort	57387	45961	52057	69065	88769	70506	54470	46064	84713	-
	Total days	480	303	392	437	228	244	338	398	306	-
	Max T	9.8	4.2	8.4	5.2	6.1	6.2	7.9	9.6	8.0	-
CKM	Min T	3.9	-0.8	2.8	0.0	0.0	1.6	1.9	3.7	2.5	-
	Mean T	6.8	1.7	5.6	2.6	3.0	3.9	4.9	6.7	5.3	-
	Tot Mort	85158	58853	69340	79306	40510	46487	60019	71098	58219	-
	Total days	524	576	361	315	294	240	279	216	134	187
	Max T	9.5	9.4	7.3	5.0	7.2	5.7	9.0	7.1	4.9	3.3
PXK	Min T	3.8	3.5	1.5	0.4	1.3	-0.6	3.8	1.4	-0.2	-1.2
	Mean T	6.6	6.5	4.5	2.7	4.2	2.6	6.4	4.2	2.4	1.0
	Tot Mort	93500	102361	63739	60352	54026	43780	51847	37852	24621	36913
	Total days	267	278	435	269	481	397	241	369	389	-
	Max T	4.5	6.7	8.9	4.9	9.9	5.8	6.4	8.5	8.6	-
SAN	Min T	-1.2	0.7	3.2	0.2	4.1	0.7	1.6	2.3	3.0	-
	Mean T	1.7	3.7	6.1	2.6	7.1	3.3	4.0	5.4	5.8	-
	Tot Mort	49316	49347	76534	52617	85053	71745	45588	65541	73249	-

281

282

283 Table 4. The number of days for each class, together with the maximum/minimum/mean
284 temperature (in °C) and the total mortality for each classification scheme in the Yorkshire
285 and Humber region. The classes showing the most oppressive (in terms of excess mortality)
286 conditions are given in bold

		CLASSES									
		1	2	3	4	5	6	7	8	9	10
	Total days	168	628	595	499	765	223	219	21	8	-
	Max T	5.8	5.1	8.9	6.0	8.3	3.3	7.2	3.7	8.8	-
ERP	Min T	1.3	0.5	3.7	0.5	2.8	-1.0	1.4	-2.5	2.9	-
	Mean T	3.5	2.8	6.3	3.3	5.6	1.1	4.3	0.6	5.9	-
	Tot Mort	30140	119958	106700	90792	144469	45468	41495	4138	1568	-
	Total days	680	472	304	439	262	210	330	209	220	-
LND	Max T	3.7	0.5	-0.6	2.5	1.5	0.3	1.8	1.7	0.4	-
	Min T	6.4	3.3	1.3	5.3	4.1	2.8	4.4	4.5	2.6	-

	Mean T	6.4	3.3	1.3	5.3	4.1	2.8	4.3	4.5	2.6	-
	Tot Mort	124020	87017	61840	81906	48071	37413	63035	39129	42296	-
KRZ	Total days	617	427	564	291	180	275	324	222	226	-
	Max T	8.8	8.9	8.2	5.8	5.9	5.6	3.9	4.1	3.9	-
	Min T	3.3	3.3	2.8	0.9	0.7	0.1	0.0	-0.2	-1.0	-
	Mean T	6.1	6.1	5.5	3.3	3.3	2.8	2.0	2.0	1.5	-
	Tot Mort	115461	77954	102629	55713	33247	49994	64012	43063	42654	-
PCT	Total days	503	554	237	433	343	298	207	294	257	-
	Max T	7.4	9.3	4.2	8.0	4.5	6.9	6.6	5.2	5.7	-
	Min T	1.9	3.8	-0.5	2.4	0.2	1.6	1.4	1.2	0.1	-
	Mean T	4.7	6.5	1.9	5.2	2.3	4.3	4.0	3.2	2.8	-
	Tot Mort	91360	101983	44293	80378	66391	54625	38984	58080	48634	-
PTT	Total days	1129	916	612	195	87	65	59	19	44	-
	Max T	8.2	7.1	5.4	4.9	4.6	3.3	4.7	4.3	4.9	-
	Min T	2.8	1.7	0.9	-0.9	0.9	-1.2	-0.4	0.3	0.4	-
	Mean T	5.5	4.4	3.1	2.0	2.8	1.1	2.2	2.3	2.7	-
	Tot Mort	207230	167469	119964	37516	16464	12831	11287	3686	8281	-
PXE	Total days	551	525	346	238	301	329	217	185	235	199
	Max T	8.2	8.5	7.5	6.8	8.0	5.1	3.8	3.9	6.3	4.8
	Min T	2.6	3.0	2.2	1.2	2.6	0.2	-0.2	-1.3	2.1	-0.2
	Mean T	5.4	5.8	4.9	4.0	5.3	2.7	1.8	1.3	4.2	2.3
	Tot Mort	103835	95659	63495	43292	54536	63661	42432	35274	46401	36142
CAP	Total days	304	236	297	388	504	391	306	245	455	-
	Max T	3.7	4.5	6.6	8.4	9.3	5.8	6.5	5.8	7.5	-
	Min T	-0.8	0.0	1.1	2.9	3.7	1.2	1.1	1.3	2.1	-
	Mean T	1.5	2.3	3.9	5.6	6.5	3.5	3.8	3.6	4.8	-
	Tot Mort	58646	46856	54418	71220	92768	71004	55650	47255	87078	-
CKM	Total days	480	303	392	437	228	244	338	398	306	-
	Max T	9.1	3.7	8.1	5.1	5.6	5.2	6.9	8.6	6.9	-
	Min T	3.5	-0.8	2.8	0.5	-0.1	0.8	1.4	2.9	1.8	-
	Mean T	6.3	1.4	5.5	2.8	2.7	3.0	4.1	5.8	4.3	-
	Tot Mort	88769	60331	71328	80227	41823	47427	61493	73613	59716	-
PXK	Total days	524	576	361	315	294	240	279	216	134	187
	Max T	8.6	8.8	6.2	4.2	6.9	5.2	8.1	6.7	4.5	2.9
	Min T	3.2	3.2	1.0	0.0	1.3	-0.6	3.2	1.6	-0.2	-1.0
	Mean T	5.9	6.0	3.6	2.1	4.1	2.3	5.6	4.1	2.1	0.9
	Tot Mort	96474	105760	65761	61338	55939	44816	53528	38712	24843	37557
SAN	Total days	267	278	435	269	481	397	241	369	389	-
	Max T	4.2	6.1	8.6	4.3	9.3	5.6	5.2	7.3	7.5	-
	Min T	-0.6	0.4	3.2	-0.2	3.6	1.0	0.8	1.7	2.2	-
	Mean T	1.8	3.3	5.9	2.1	6.4	3.3	3.0	4.5	4.9	-
	Tot Mort	50373	50491	79640	53780	88491	72329	46762	67780	75083	-

287

288

289 Table 5. The number of days for each class, together with the maximum/minimum/mean
290 temperature (in °C) and the total mortality for each classification scheme in the Northeast
291 region. The classes showing the most oppressive (in terms of excess mortality) conditions
292 are given in bold

		CLASSES									
		1	2	3	4	5	6	7	8	9	10
ERP	Total days	168	628	595	499	765	223	219	21	8	-
	Max T	7.1	6.3	9.9	6.9	9.0	4.9	7.8	5.1	9.7	-
	Min T	2.7	2.0	4.6	1.7	3.8	0.9	2.2	-0.9	4.1	-
	Mean T	4.9	4.2	7.2	4.3	6.4	2.9	5.0	2.1	6.9	-
	Tot Mort	31723	121646	110930	93821	148527	46415	43598	4411	1486	-
LND	Total days	680	472	304	439	262	210	330	209	220	-
	Max T	9.7	7.0	5.1	8.9	7.6	6.2	7.8	8.4	6.2	-
	Min T	4.3	1.8	1.4	3.6	2.5	1.4	2.9	2.9	2.3	-
	Mean T	7.0	4.4	3.3	6.2	5.0	3.8	5.3	5.7	4.2	-
	Tot Mort	128481	90218	62648	84060	50113	38281	64093	40887	43774	-
KRZ	Total days	617	427	564	291	180	275	324	222	226	-
	Max T	9.5	9.6	8.9	6.7	6.8	6.6	5.6	5.7	5.3	-
	Min T	4.0	4.2	3.7	2.2	2.2	1.5	1.8	1.6	0.8	-
	Mean T	6.7	6.9	6.3	4.5	4.5	4.0	3.7	3.6	3.0	-
	Tot Mort	118771	80666	106411	57162	34675	52260	64751	44071	43788	-
PCT	Total days	503	554	237	433	343	298	207	294	257	-
	Max T	8.2	10.0	5.7	8.9	5.9	7.3	7.7	6.5	6.7	-
	Min T	2.9	4.5	1.3	3.6	1.8	2.4	2.7	2.9	1.3	-
	Mean T	5.6	7.2	3.4	6.3	3.9	4.8	5.2	4.7	4.0	-
	Tot Mort	94384	105703	45399	82492	68026	56254	39882	59317	51099	-
PTT	Total days	1129	916	612	195	87	65	59	19	44	-
	Max T	8.9	8.0	6.7	6.1	6.5	5.1	6.1	5.8	6.1	-
	Min T	3.7	2.8	2.6	0.7	2.8	0.9	1.3	2.3	1.3	-
	Mean T	6.3	5.4	4.6	3.4	4.7	3.0	3.7	4.1	3.7	-
	Tot Mort	214404	173334	121906	39016	16864	13065	11724	3781	8463	-
PXE	Total days	551	525	346	238	301	329	217	185	235	199
	Max T	9.2	9.0	8.5	7.5	8.8	6.4	5.6	5.3	7.3	5.8
	Min T	3.7	3.8	3.3	2.0	3.5	2.1	1.8	0.5	3.2	1.1
	Mean T	6.5	6.4	5.9	4.7	6.2	4.2	3.7	2.9	5.3	3.4
	Tot Mort	106007	99439	65893	44744	57061	65197	43109	36551	47607	36946
CAP	Total days	304	236	297	388	504	391	306	245	455	-
	Max T	5.3	5.8	7.7	9.3	9.9	7.5	7.3	6.8	7.9	-
	Min T	1.1	1.5	2.4	3.9	4.3	2.8	2.2	2.7	2.9	-
	Mean T	3.2	3.7	5.0	6.6	7.1	5.1	4.8	4.8	5.5	-
	Tot Mort	60164	47475	56192	73555	94323	73811	57479	48893	90664	-
CKM	Total days	480	303	392	437	228	244	338	398	306	-
	Max T	9.9	5.2	9.1	6.8	6.5	6.3	7.6	8.9	7.5	-
	Min T	4.3	1.0	3.9	2.3	1.3	2.4	2.4	3.6	2.8	-
	Mean T	7.1	3.1	6.5	4.5	3.9	4.4	5.0	6.3	5.2	-
	Tot Mort	90601	61076	73705	83262	43222	48212	63799	76486	62193	-
P XK	Total days	524	576	361	315	294	240	279	216	134	187
	Max T	9.4	9.6	6.8	5.8	8.0	6.2	8.7	7.7	5.7	4.8
	Min T	3.8	4.2	1.9	1.8	2.7	0.9	4.1	2.9	1.3	1.1
	Mean T	6.6	6.9	4.4	3.8	5.4	3.6	6.4	5.4	3.5	2.9
	Tot Mort	100529	108783	67380	62707	57836	46159	54956	40145	25587	38473
SAN	Total days	267	278	435	269	481	397	241	369	389	-
	Max T	5.7	6.9	9.5	5.6	9.9	7.3	6.4	8.0	7.9	-

Min T	1.2	1.8	4.1	1.5	4.3	2.7	2.3	2.7	3.1	-
Mean T	3.4	4.3	6.8	3.6	7.0	5.0	4.3	5.4	5.5	-
Tot Mort	52010	52670	81556	54696	90012	75071	47937	70319	78283	-

293


294 3.2 Analysis of the atmospheric patterns

295 In order to study the climatology of the most oppressive, as defined above, classes for the
 296 five regions studied, centroid maps of the surface atmospheric pressure regimes are given in
 297 Figure 4. It appears that all Figure 4 maps more-or-less depict the same circulation patterns,
 298 i.e. the most dangerous class, in terms of excess winter mortality for each classification
 299 scheme is characterized by weak (rather shallow) low atmospheric pressure systems located
 300 west (and sometimes southwest, for example see ERP class 6 and LND class 3) of the British
 301 Isles, whereas anti-cyclonic conditions prevail over Scandinavia and west continental Europe
 302 for almost all cases. It is also noteworthy that the semi-permanent Icelandic Low seems to
 303 be absent in all 10 classification schemes.

304 This type of weather is known as the Easterly type and is generally characterized by
 305 anticyclones over Scandinavia (sometimes extending towards Iceland) and depressions
 306 circulating over the western North Atlantic, as well as the Bay of Biscay, where the Azores
 307 Anticyclone is generally absent (Lamb, 1950). The aforementioned Easterly type of weather
 308 is associated with low temperatures in autumn, winter and spring and sometimes extremely
 309 low temperatures with occasional snow in southern districts, snow or sleet showers in
 310 eastern and northeastern districts, and dry conditions in western regions (Lamb, 1950). In
 311 terms of air masses' movement, the Easterly type of weather is associated with advection of
 312 air originating from continental Europe. These flows are characterized by subsidence of
 313 several hundred hPa before they land (Walsh et al., 2001) and they are responsible for the
 314 low temperatures observed in England, as opposed to relatively warm and humid air
 315 originating from the Atlantic. Walsh et al. (2001) and Cattiaux et al. (2013) state that this
 316 atmospheric pattern is related to a negative phase of the North Atlantic Oscillation, as well
 317 as to positive pressure anomalies over the Arctic.


318

MSLP(year) for ERP09_YR_S04_SP_D04 type #06




320 (a)

MSLP(year) for LND09_YR_S04_SP_D04 type #03




(b)

MSLP(year) for KRZ09_YR_S04_SP_D04 type #07




322 (c)

MSLP(year) for PCT09_YR_S04_SP_D04 type #08




(d)

MSLP(year) for PTT09_YR_S04_SP_D04 type #03



324 (e)

MSLP(year) for PXE09_YR_S04_SP_D04 type #09



(f)




Figure 4. Centroid maps of surface atmospheric pressure for the most oppressive conditions, as described by ERP class 6 (a), LND class 3 (b), KRZ class 7 (c), PCT class 8 (d), PTT class 3 (e), PXE class 9 (f), CAP class 2 (g), CKM class 2 (h), PXK class 4 (i), SAN class 4 (j).

According to Dimitriou et al. (2016), easterly flows in England are very often associated with unfavorable conditions for public health, as the advection of very cold continental air from northern/eastern Europe that continues for several days, results in low temperatures and excess mortality. This type of weather is very often associated with a blocking pattern over Western Europe or increased pressure over Eastern/Northern Europe. Specifically for the northwest region, Dimitriou et al. (2016) found that easterly short-range flows were linked to a 5.3% increase in winter mortality.

4. Conclusions

The objective of the present work has been to study the link between atmospheric patterns and cold-related mortality at the daily time-scale in England, in order to shed light on the climatology of health outcomes. In doing so, 10 different classification methods provided by the COST-733 tool were used. The use of the specific classification techniques brought a new perspective to the understanding of the climatological associations between mortality and winter weather, and appears to be a valuable addition to the suite of tools available for climate/health data analysis.

Our results showed that the most unfavorable conditions for public health in the 5 regions of England were associated with Easterly weather, which is known to favor advection of cold air originating from continental Europe. The fact that little-to-no variation among the 5 regions was observed, when grouping the days to form the most oppressive classes for public health, suggested a spatially homogeneous response of the population to the specific atmospheric patterns identified.

Not surprisingly, in most of the cases examined through the different classification schemes, excess mortality was linked to the lowest daily minimum/maximum temperatures in the 5 regions, in agreement with previous researchers. Nevertheless in a number of cases high mortality rates were associated with relatively higher temperatures, probably due to rapidly changing weather. This finding is indicative of the complexity of cold-related health outcomes and highlights the role of synoptic climatology on confounding the relationship between temperature and mortality.

On the whole, the results provided here show that although cold-related health outcomes can be fatal, they can also be predictable and preventable (Ghosh et al., 2014), as policy-makers can be informed appropriately and design intervention strategies towards allocating resources and reducing the adverse health effects of cold weather. In any case further analysis is needed to clarify how the various climatic elements can increase population vulnerability. Additionally, further analysis with more recent data is needed, where due consideration should be given to how to control for regional variation due to socioeconomic differences.

References

- Allen, M. J., and C. C. Lee, 2014: Investigating high mortality during the cold season: Mapping mean weather patterns of temperature and pressure. *Theor. Appl. Climatol.*, 118, 419–428, doi:10.1007/s00704-013-1075-x.
- Allen, M.J. and S.C Sheridan, 2014: High-mortality days during the winter season: comparing meteorological conditions across 5 US cities. *Int J Biometeorol*, 58(2), 217-252, doi: 10.1007/s00484-013-0640-4.
- Analitis, A., and Coauthors, 2008: Effects of cold weather on mortality: Results from 15 European cities within the PHEWE project. *Amer. J. Epidemiol.*, 168, 1397–1408, doi:10.1093/aje/kwn266.
- Anderson, B. G., and M. L. Bell, 2009: Weather-related mortality: How heat, cold, and heat waves affect mortality in the United States. *Epidemiology*, 20, 205–213, doi:10.1097/EDE.0b013e318190ee08.

- Astrom, D. O., B. Forsberg, S. Edvinsson, and J. Rocklov, 2013: Acute fatal effects of short-lasting extreme temperatures in Stockholm, Sweden: Evidence across a century of change. *Epidemiology*, 24, 820–829, doi:10.1097/01.ede.0000434530.62353.0b.
- Bhaskaran, K., S. Hajat, A. Haines, E. Herrett, P. Wilkinson, and L. Smeeth, 2010: Short term effects of temperature on risk of myocardial infarction in England and Wales: Time series regression analysis of the Myocardial Ischaemia National Audit Project (MINAP) registry. *BMJ*, 341, C3823, doi:10.1136/bmj.c3823.
- Basu, R., 2009. High ambient temperature and mortality: a review of epidemiologic studies from 2001 to 2008. *Environ. Heal.* 8, 40.
- Basu, R., Samet, J., 2002. Relation between elevated ambient temperature and mortality: a review of the epidemiologic evidence. *Epidemiol. Rev.* 24, 190–202.
- Basu, R., Feng, W.Y., Ostro, B.D., 2008. Characterizing temperature and mortality in nine California counties. *Epidemiology* 19, 138–145.
- Carson, C., S. Hajat, B. Armstrong, and P. Wilkinson, 2006: Declining vulnerability to temperature related mortality in London over the 20th century. *Amer. J. Epidemiol.*, 164, 77–84, doi:10.1093/aje/kwj147.
- Cattiaux, J., H. Douville, A. Ribes, F. Chauvin, and C. Plante, 2013: Towards a better understanding of changes in wintertime cold extremes over Europe: A pilot study with CNRM and IPSL atmospheric models. *Climate Dyn.*, 40, 2433–2445, doi:10.1007/s00382-012-1436-7.
- Curriero, F. C., K. S. Heiner, J. M. Samet, S. L. Zeger, L. Strug, and J. A. Patz, 2002: Temperature and mortality in 11 cities of the eastern United States. *Amer. J. Epidemiol.*, 155, 80–87, doi:10.1093/aje/155.1.80.
- de'Donato, F. K., M. Leone, D. Noce, M. Davoli, and P. Michelozzi, 2013: The impact of the February 2012 cold spell on health in Italy using surveillance data. *PLoS One*, 8, e61720, doi:10.1371/journal.pone.0061720.
- Dimitriou, K., McGregor, G., Kassomenos, P., Paschalidou, A., (2016): Exploring winter mortality variability in five regions of England, using back trajectory analysis. *Earth Interactions*, 20 (1), doi:10.1175/EI-D-15-0012.1.
- Donaldson, G. C., and W. R. Keatinge, 2003: Cold related mortality in England and Wales; influence of social class in working and retired age groups. *J. Epidemiol. Commun. Health*, 57, 790–791, doi:10.1136/jech.57.10.790.
- Donaldson, G.C., Keatinge W.R., and S. Nayha, 2003: Changes in summer temperature and heat-related mortality since 1971 in North Carolina, south Finland, and southeast England. *Environ. Res.*, 91, 1–7, doi:10.1016/S0013-9351(02)00002-6.
- Duckstein, L., Bardossy, A., Bogardi, I., 1993. Linkage between the occurrence of daily atmospheric circulation patterns and floods: an Arizona case study. *J. Hydrol.* 143, 413–428.
- Erpicum, M., Mabilie, G., Fettweis, X., 2008. Automatic synoptic weather circulation types classification based on the 850 hPa geopotential height. Abstracts COST 733 Midterm Conference, Advances in Weather and Circulation Type Classifications & Applications 22–25 October 2008, p. 33 Krakow, Poland.
- Ferreira Braga, A. L., A. Luis, A. Zanobetti, and J. Schwartz, 2001: The time course of weather related deaths. *Epidemiology*, 12, 662–667, doi:10.1097/00001648-200111000-00014.
- Gasparini A, B. Armstrong, and M. G. Kenward, 2010: Distributed lag nonlinear models. *Stat. Med.*, 29, 2224–2234, doi:10.1002/sim.3940.
- Gasparrini, A., and M. Leone, 2014: Attributable risk from distributed lag models. *BMC Med. Res. Methodol.*, 14, 55, doi:10.1186/1471-2288-14-55.

- Ghosh, A., C. Carmichael, A.J. Elliot, H.K. Green, V. Murray, C. Petrokofsky, 2014: The Cold Weather Plan evaluation: an example of pragmatic evidence-based policy making? *Public Health*, 128(7), 619-627.
- Gomez-Acebo, I., T. Dierssen-Sotos, and J. Llorca, 2010: Effect of cold temperatures on mortality in Cantabria (northern Spain): A case-crossover study. *Public Health*, 124, 398–403, doi:10.1016/j.puhe.2010.03.025.
- Gosling, S. N., J. A. Lowe, G. R. McGregor, M. Pelling, and B. D. Malamud, 2009: Associations between elevated atmospheric temperature and human mortality: A critical review of the literature. *Climatic Change*, 92, 299–341, doi:10.1007/s10584-008-9441-x.
- Guo, Y., S. Li, Y. Zhang, B. Armstrong, J. J. K. Jaakkola, S. Tong, and X. Pan, 2013: Extremely cold and hot temperatures increase the risk of ischaemic heart disease mortality: Epidemiological evidence from China. *Heart*, 99, 195–203, doi:10.1136/heartjnl-2012-302518.
- Guo, Y., Gasparrini, A., Armstrong, B., Li, S., Tawatsupa, B., Tobias, A., et al., 2014. Global variation in the effects of ambient temperature on mortality: a systematic effect. *Epidemiology* 25, 781–789.
- Hajat, S, R. S. Kovats, and K. Lachowycz, 2007: Heat-related and cold-related deaths in England and Wales: Who is at risk? *Occup. Environ. Med.*, 64, 93–100, doi:10.1136/oem.2006.029017.
- Hajat, S., and S. Kovats, 2014: A note of caution about the excess winter deaths measure. *Nat. Climate Change*, 4, 647, doi:10.1038/nclimate2302.
- Hajat, S., Vardoulakis, S., Heaviside, C., Eggen, B., 2014. Climate change effects on human health: projections of temperature-related mortality for the UK during the 2020s, 2050s and 2080s. *J. Epidemiol. Community Health* 68, 641–648.
- Healy, J. D., 2003: Excess winter mortality in Europe: A cross country analysis identifying key risk factors. *J. Epidemiol. Commun. Health*, 57, 784–789, doi:10.1136/jech.57.10.784.
- Heaviside, C., Tsangari, H., Paschalidou, A., Vardoulakis, S., Kassomenos, P., Georgiou, K.E., Yamasaki E.N., (2016): Heat-related mortality in Cyprus for current and future climate scenarios. *The Science of the Total Environment*, 569-570, 627-633.
- Huang, C., Barnett, A.G., Wang, X., Vaneckova, P., FitzGerald, G., Tong, S., 2011. Projecting future heat-related mortality under climate change scenarios: a systematic review. *Environ. Health Perspect.* 119, 1681–1690.
- Keatinge, W.R., Donaldson, G.C., Cordoli, E., Martinelli, M., Kunst, A.E., Mackenbach, J.P., et al., 2000. Heat related mortality in warm and cold regions of Europe: observational study. *BMJ* 321, 670–673.
- Keatinge, W. R., 2002: Winter mortality and its causes. *Int. J. Circumpolar Health*, 61, 292–99, doi:10.3402/ijch.v61i4.17477.
- Kovats, S., and T. Kosatsky, 2009: Similarities and contrasts in the social and environmental determinants of weather-related mortality: Heat versus cold. *Epidemiology*, 20, S235, doi:10.1097/01.ede.0000362789.62314.f0.
- Lamb, H. H., 1950: Types and spells of weather around the year in the British Isles: Annual trends, seasonal structure of the year, singularities. *Quarterly Journal of the Royal Meteorological Society*, 76 (330), 393-429.
- Lund, I.A., 1963. Map-pattern classification by statistical methods. *J. Appl. Meteorol.* 2, 56–65.
- McGregor G.R, 2005: Winter North Atlantic Oscillation, temperature and ischaemic heart disease mortality in three English counties. *Int. J. Biometeor.*, 49, 197–204, doi:10.1007/s00484-004-0221-7.

- Montero, J. C., I. J. Miron, J. J. Criado-Alvarez, C. Linares, and J. Diaz, 2010: Mortality from cold waves in Castile—La Mancha, Spain. *Sci. Total Environ.*, 408, 5768–5774, doi:10.1016/j.scitotenv.2010.07.086.
- Morabito, M., and Coauthors, 2012: Air temperature-related human health outcomes: Current impact and estimations of future risks in central Italy. *Sci. Total Environ.*, 441, 28–40, doi:10.1016/j.scitotenv.2012.09.056.
- Ma, W., C. Yang, C. Chu, T. Li, J. Tan, and H. Kan, 2013: The impact of the 2008 cold spell on mortality in Shanghai, China. *Int. J. Biometeor.*, 57, 179–184, doi:10.1007/s00484-012-0545-7.
- Paschalidou, A.K., Kassomenos P.A. (2016): What are the most fire-dangerous atmospheric circulations in the Eastern-Mediterranean? Analysis of the synoptic wildfire climatology. *Science of the Total Environment*, 539, 536–545.
- Philipp, A., Bartholy, J., Beck, C., Erpicum, M., Esteban, P., Fettweis, R., Huth, R., James, P., Jourdain, S., Kreienkamp, F., Krennert, T., Lykoudis, S., Michaelides, S., Pianko, K., Post, P., Rasilla, A.D., Schiemann, R., Spekat, A., Tymvios, F.S., 2010. Cost733cat — a database of weather and circulation type classifications. *Phys. Chem. Earth* 35, 360–373.
- Philipp, A., Beck, C., Huth, R., Jacobeit, J., 2016. Development and comparison of circulation type classifications using the COST 733 dataset and software. *Int. J. Climatol.* 36: 2673–2691, doi:10.1002/joc.3920.
- Scarborough, P., S. Allender, M. Rayner, and M. Goldacre, 2012: Contribution of climate and air pollution to variation in coronary heart disease mortality rates in England. *PLoS One*, 7, e32787, doi:10.1371/journal.pone.0032787.
- Tsangari, H., Paschalidou, A., Vardoulakis, S., Heaviside, C., Konsoula, Z., Christou, S., Georgiou, K.E., Ioannou, K., Mesimeris, T., Kleanthous, S., Pashiardis, S., Pavlou, P., Kassomenos, P., Yamasaki E.N. (2016): Human mortality in Cyprus: the role of temperature and particulate air pollution. *Regional Environmental Change*, 16(7), 1905–1913.
- Urban, A., H. Davidkovova, and J. Kysely, 2014: Heat- and cold-stress effects on cardiovascular mortality and morbidity among urban and rural populations in the Czech Republic. *Int. J. Biometeor.*, 58, 1057–1068, doi:10.1007/s00484-013-0693-4.
- Vardoulakis, S., Dear, K., Hajat, S., Heaviside, C., Eggen, B., McMichael, A.J., 2014: Comparative Assessment of the Effects of Climate Change on Heat- and Cold-Related Mortality in the United Kingdom and Australia, *Environmental Health Perspectives*, 122, 1285–1292.
- Vasconcelos, J., E. Freire, R. Almendra, G. L. Silva, and P. Santana, 2013: The impact of winter cold weather on acute myocardial infarctions in Portugal. *Environ. Pollut.*, 183, 14–18, doi:10.1016/j.envpol.2013.01.037.
- Wang, C., R. Chen, X. Kuang, X. Duan, and H. Kan, 2014: Temperature and daily mortality in Suzhou, China: A time series analysis. *Sci. Total Environ.*, 466–467, 985–990, doi:10.1016/j.scitotenv.2013.08.011.
- Wang Y., Shi L., Zanobetti A., Schwartz J.D., 2016: Estimating and projecting the effect of cold waves on mortality in 209 US cities, *Environment International* 94, 141–149.
- Walsh, J. E., A. S. Phillips, D. H. Portis, and W. L. Chapman, 2001: Extreme cold outbreaks in the United States and Europe, 1948–99. *J. Climate*, 14, 2642–2658, doi:10.1175/1520-0442(2001)014<2642:ECOITU.2.0.CO;2.
- Xie, H. Y., and Coauthors, 2013: Short-term effects of the 2008 cold spell on mortality in three subtropical cities in Guangdong Province, China. *Environ. Health Perspect.*, 121, 210–216, doi:10.1289/ehp.1104541.

534 Xu, W., and Coauthors, 2013: Thermal stress associated mortality risk and effect
535 modification by sex and obesity in an elderly cohort of Chinese in Hong Kong.
536 *Environ. Pollut.*, 178, 288–293, doi:10.1016/j.envpol.2013.03.020.

537 Yu, W., Mengersen, K., Wang, X., Ye, X., Guo, Y., Pan, X., Tong, S., 2012. Daily average
538 temperature and mortality among the elderly: a meta-analysis and systematic
539 review of epidemiological evidence. *Int. J. Biometeorol.* 56, 569–581.

540 Zeka, A., S. Browne, H. McAvoy, and P. Goodman, 2014: The association of cold weather and
541 all cause and cause-specific mortality in the island of Ireland between 1984 and
542 2007. *Environ. Health*, 13, 104, doi:10.1186/1476-069X-13-104.

543 Zhou, M. G., and Coauthors, 2014: Health impact of the 2008 cold spell on mortality in
544 subtropical China: The Climate and Health Impact National Assessment Study
545 (CHINAs). *Environ. Health*, 13, 60, doi:10.1186/1476-069X-13-60.